## Physics

Question	Maximum Mark	Mark Awarded
#1	6	
#2	6	
#3	8	
#4	9	
#5	9	
#6	11	
#7	20	
Total	69	



Disclaimer: The questions in this revision paper have all been taken from actual examinations that have taken place. Whilst the questions are the property of Eduqas, this revision paper was created using an online tool and Eduqas take no responsibility for the content within it.

7	7.	(a)	Explain briefly what is meant by conservation of mass-energy.	[2]
		(b)	It is suggested that a collision between two protons, each of kinetic energy 3 GeV producthe following interaction:	es
			$p + p \longrightarrow 5p + 3\bar{p} + n + \bar{n} + 2\pi^{+} + 2\pi^{-} + 4\nu_{e}$	
			Determine which, if any, of the conservation laws are violated (the rest mass-energy of proton or a neutron ≈ 1 GeV).	f a [4]

Question taken from Eduqas examination paper 842103, June 2018

Discuss the make-up and properties of the following particles $e^-, e^+, n, \overline{p}, \pi^-$ .	[6 QER]

Question taken from Eduqas examination paper 842103, November 2020

	_
++	Z
-++	

6.	(a)	The anti-Δ++ is an anti-baryon and a first-generation particle which has a charge of –2 Explain why the only possible quark make-up of the anti-Δ++ is ūūū.
	(b)	The anti- $\Delta^{++}$ has a lifetime of approximately $6\times 10^{-24} s$ and decays into a $\pi$ meson ar another anti-baryon. Deduce the quark make-up of the $\pi$ meson and the anti-baryon ar name them.
	(c)	State which force is responsible for the decay of the anti- $\Delta^{++}$ into a $\pi$ meson and an baryon, giving a reason for your answer.
	(d)	In 2011, a highly respected international research collaboration reported that they had measured neutrinos travelling at speeds greater than that of light. This report was met light caution from the scientific community and then the result was disproved. Explain brief why the result was met with caution and how the results might have been disproved. [

Question taken from Eduqas examination paper 842103, June 2019

#4

## theonlinephysicstutor.com

When colliding two protons at very high energy the following interaction has been observed at the large hadron collider in CERN.

$p + p \rightarrow {}_{1}^{2}H + e^{+} + 3$	p.	+ 1	) –	→ <sup>2</sup>	Η	+	e <sup>+</sup>	+	5
---	----	-----	-----	----------------	---	---	----------------	---	---

(a)	Use the conservation of lepton number and conservation of charge to identify the pa	ticle [3]
(b)	State which force is responsible for this interaction. Give your reasoning.	[2]
(c)	State which of the above particles can be classed as:  (i) a baryon;	[2]
(d)	(ii) an antiparticle.  At present, countries in Europe contribute to funding research at CERN. Evaluate who or not the money could be better spent on humanitarian aid.	ether [2]

Question taken from Eduqas examination paper 842001, June 2017

0	7.	(a)	Quarks and electron neutrinos are fundamental particles whereas protons and neut are not. Explain this statement.	trons [2]
		(b)	(i) State why it is that electron neutrinos are very difficult to detect.	[2]
			(ii) Electron neutrinos can be detected when they interact with deuterium nuclei, that are present in heavy water. The following reaction is observed.	, <sup>2</sup> H,
			$v_{\rm e}$ + $_{\rm 1}^{\rm 2}{ m H}$ $ ightarrow$ p + p + x Identify particle x and justify your answer using the conservation laws.	[3]
		(c)	When a positron and an electron meet they annihilate to produce two gamma ray photo $e^+ + e^- \to \gamma + \gamma$	tons.
			State which force is responsible for this interaction, giving your reasoning.	[2]

Question taken from Eduqas examination paper 842001, June 2019

5.	(a)	Subatomic particles can be classified as either hadrons or leptons. Giving examples of each, fully describe the differences and similarities between these two groups. Include the types of interaction they undergo and how one of the groups can be further sub divided.  [6 QER]

atom	n. The following interaction occurs:	
	p + e <sup>−</sup> → n + x	
(i)	Identify the particle, $x$ , explaining how you used the relevant conservation laws	s.[3]
(ii)	Describe the change in quark flavour during electron capture.	[1]
(iii)	State and explain which interaction is responsible for this decay.	[1]

Electron capture occurs when an electron interacts with a proton in the nucleus of an

Question taken from Eduqas examination paper 842001, June 2018

Λ	arbon-14 nucleus decays as shown:	theonlinephys
A cai	${}^{16}C \longrightarrow {}^{17}N + e^- + \overline{\nu_e}$	
(i)	Show how charge, baryon number and le	
	decay.	
(ii)	Give two reasons why this must be a weak n	nuclear force interaction.
The o	decay constant of carbon-14 is $3.83 \times 10^{-12}$ s	-1
(i)	Calculate its half-life in years.	-
(ii)	The natural ratio of carbon-14 to carbon-12	is 1.00 × 10 <sup>-12</sup> i.e.
	number of <sup>14</sup> C nuclei	
	1000000000000000000000000000000000000	× 10 <sup>-12</sup>
	Calculate the activity of 12 g of naturally occ	surring carbon
	Calculate the activity of 12 g of flaturally occ	diffig carbon.
(iii)		
	decayed but the carbon-12 all remains. The old tree has dropped to 0.34 × 10 <sup>-12</sup> . Calcu	late the age of the old tree.
Carbo	oon-11 ( $^{11}_{ m C}$ C) is proton rich and undergoes positr Complete the following decay equation for cart	on decay to a stable isotope of b
requir	ire analysis of lepton number, baryon number a	and charge.
	11 C →	
	0	
Some	he 14 March 2013, the discovery of the Higgs been physicists were convinced that they had dived that there are many different types of Higg	discovered the Higgs boson, o
	just another particle and not the Higgs bosor ded which, if any, of these claims is correct.	n. Explain how it may or may no
was ji		
was ji		

Question taken from Eduqas examination paper 842103, June 2017