


Physics

Question	Maximum Mark	Mark Awarded
#1	20	
#2	20	
#3	20	
#4	20	
Total	80	

created with
 **Question Bank**
Part of WJEC

Disclaimer: The questions in this revision paper have all been taken from actual examinations that have taken place. Whilst the questions are the property of Eduqas, this revision paper was created using an online tool and Eduqas take no responsibility for the content within it.

#1

13. The following photograph shows a cyclist crossing the finish line at the end of a race.



(a) State why the net moment acting on the cyclist is zero. [1]

.....

.....

.....

(b) (i) State what is meant by the moment of inertia of an object. [2]

.....

.....

.....

(ii) A diver of mass 60 kg and height 1.68 m has an angular momentum of $92.1 \text{ kg m}^2 \text{ s}^{-1}$ at the start of her dive. Her moment of inertia, I , at the start of the dive is

$$I = \frac{1}{12} mh^2$$

where m is her mass and h is her height. Calculate her angular velocity. [3]

.....

.....

.....

.....

.....

- (iii) During the dive, the diver tucks in her arms and legs and reduces the moment of inertia to 2.7 kg m^2 . Calculate the final angular velocity of the diver. [3]

.....

.....

.....

.....

.....

- (c) The wheel of a Formula 1 car has a moment of inertia of 1.10 kg m^2 . As the car approaches a corner and brakes, its angular velocity decreases from 220 rad s^{-1} to 170 rad s^{-1} in a time of 0.310 s .

- (i) Calculate the resultant torque on the wheel of the car during the braking process. [3]

.....

.....

.....

.....

.....

- (ii) Determine the total rotational kinetic energy lost by the wheels of the car during the above braking process assuming all the wheels have the same moment of inertia. [3]

.....

.....

.....

.....

.....

- (d) During a Grand Prix a driver loses control of the car when approaching a bend and crashes but escapes with minor injuries. The speed reduces from 213 km hr^{-1} to zero in a time of 0.651 s . The mass of the car is 640 kg and the driver's mass is 70 kg . Use the given data to evaluate why Grand Prix race circuits have large areas of grass or loose stone chippings around certain corners. [5]

.....

.....

.....

.....

.....

.....

.....


.....

.....

Question taken from Eduqas examination paper 842103, June 2018

#2

13. (a) The diagram shows the forces acting on an athlete's arm as she holds a discus. Calculate the magnitude of the tension, T , provided by the deltoid muscle. [3]



.....

.....

.....

.....

- (b) (i) Define angular acceleration. [1]

.....

.....

.....

- (ii) When thrown, the discus experiences an angular acceleration. It accelerates from rest to 2.3 revolutions per second in a time of 0.27s. Calculate the angular acceleration of the discus. [2]

.....

.....

.....

.....


- (iii) Calculate the mean torque applied to the discus of mass 2.0 kg and radius 11 cm. The moment of inertia is given by the equation $I = \frac{mr^2}{2}$. [3]

.....

.....

.....

- (c) (i) Calculate the maximum height attained by the discus if it is thrown with a velocity of 24 m s^{-1} and an angle of 38° from a height of 1.2 m . *Ignore the effects of air on the motion of the discus.* [4]



.....

.....


.....

.....

.....

.....

- (ii) Taking the effects of air into account, evaluate whether the horizontal distance travelled by the discus will increase, decrease or remain approximately the same. The diagram shows the speed of air relative to the upper and lower surfaces of the discus. The density of air = 1.3 kg m^{-3} . [5]



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) A wind tunnel is used to examine the motion of the discus in a wind speed of 20 ms^{-1} . Calculate the factor by which the drag force increases if the speed of the wind is increased to 30 ms^{-1} and all other factors are kept constant. [2]

.....

.....


.....

.....

Question taken from Eduqas examination paper 842103, June 2019

#3

- (a) (i) At the start of a tennis game, a player serves the ball with an initial velocity of 44 m s^{-1} at an angle of 7° to the horizontal as shown below. The maximum horizontal distance for the ball to stay in play is 18.29 m . If the ball remains in the air for a time of 0.41 s , determine if the ball lands in play from the serve. *Ignore the effects of air resistance for this part of the question.* [3]




- (ii) The tennis ball has a mass of 0.056 kg and is momentarily at rest before being hit by the racquet. Determine the mean force exerted by the racquet on the ball if they remain in contact for a time of 6.0 ms . [2]

- (b) (i) The coefficient of restitution between the ball and the floor is 0.74 . Explain what this statement means. [2]

- (ii) Determine the **second** bounce height of a tennis ball if the ball is dropped from a height of 1.95 m (the coefficient of restitution between the ball and the floor is 0.74). [3]

- (c) (i) During the game, the player plays a shot and applies spin to the ball. Explain how the ball will travel through the air by discussing the forces acting on the ball. Label the forces and their directions on the diagram provided. [4]



- (ii) The ball is hit with a velocity of 16.4 m s^{-1} and spins at a rate of 3500 revolutions per minute. Determine the **total** kinetic energy of the ball if the diameter of the ball is 7.0 cm and its mass is 0.056 kg .
Note: a tennis ball can be considered to be a thin spherical shell. [4]

- (iii) Determine the drag force acting on the ball if the drag coefficient for a tennis ball is 0.53 and the density of air is 1.2 kg m^{-3} . [2]

Question taken from Eduqas examination paper 842103, June 2017

#4

- (a) (i) Explain what is meant by the term *moment of inertia* of an object. [2]

.....

.....

- (ii) Calculate the moment of inertia of a cricket ball which has a rotational kinetic energy of 1.47 J if it is spinning at a rate of 30 revolutions per second. [3]

.....

.....

.....

- (b) (i) The batsman hits the ball with an initial velocity of 25 m s^{-1} at an angle of 30° to the horizontal. A fielder standing 5.6 m away from the batsman can catch a ball 2.4 m above the ground. Evaluate whether the ball can be caught by the fielder. Assume that air resistance can be ignored and that the ball is hit from ground level. [5]



.....

.....

.....

.....

.....

- (ii) Explain why a fielder will move his hands in the direction of motion of the cricket ball when catching. [2]

.....

.....

- (iii) The coefficient of restitution between the pitch and the ball is 0.37. Determine the bounce height if the ball falls from a height of 2.35 m. [2]

.....

.....

- (c) For this part of the question, the interactions between the ball and the air need to be taken into account.

- (i) Explain why a spinning cricket ball will change direction when moving through the air. Your answer should include the forces acting on the ball during the flight and a diagram may be included. [3]

.....

.....

.....

.....

- (ii) Determine the drag force acting on a cricket ball of radius 3.6 cm during flight if the speed of the ball is 24.3 m s^{-1} and its drag coefficient is 0.76. Density of air = 1.3 kg m^{-3} . [3]

.....

.....

.....

Question taken from Eduqas examination paper 842103, November 2020